

Complete Genome Sequences of Six Strains of the Genus *Methylobacterium*

Christopher J. Marx, Françoise Bringel, Ludmila Chistoserdova, Lionel Moulin, Muhammad Farhan UI Haque, Darrell E. Fleischman, Christelle Gruffaz, Philippe Jourand, Claudia Knief, Ming-Chun Lee, Emilie E. L. Muller, Thierry Nadalig, Rémi Peyraud, Sandro Roselli, Lina Russ, Lynne A. Goodwin, Natalia Ivanova, Nikos Kyrpides, Aurélie Lajus, Miriam L. Land, Claudine Médigue, Natalia Mikhailova, Matt Nolan, Tanja Woyke, Sergey Stolyar, Julia A. Vorholt and Stéphane Vuilleumier
J. Bacteriol. 2012, 194(17):4746. DOI: 10.1128/JB.01009-12.

Updated information and services can be found at:
<http://jb.asm.org/content/194/17/4746>

	<i>These include:</i>
REFERENCES	This article cites 17 articles, 6 of which can be accessed free at: http://jb.asm.org/content/194/17/4746#ref-list-1
CONTENT ALERTS	Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), more»

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Complete Genome Sequences of Six Strains of the Genus *Methylobacterium*

Christopher J. Marx,^{a,b} Françoise Bringel,^c Ludmila Chistoserdova,^d Lionel Moulin,^e Muhammad Farhan Ul Haque,^c Darrell E. Fleischman,^f Christelle Gruffaz,^c Philippe Jourand,^e Claudia Knief,^g Ming-Chun Lee,^a Emilie E. L. Muller,^c Thierry Nadalig,^c Rémi Peyraud,^g Sandro Roselli,^c Lina Russ,^g Lynne A. Goodwin,^{h,i} Natalia Ivanova,^{i,j} Nikos Kyrpides,ⁱ Aurélie Lajus,^k Miriam L. Land,^{i,l} Claudine Médigue,^k Natalia Mikhailova,^{i,j} Matt Nolan,ⁱ Tanja Woyke,^{i,j} Sergey Stoliar,^m Julia A. Vorholt,^g and Stéphane Vuilleumier^c

Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, Massachusetts, USA^a; Faculty of Arts and Sciences Center for Systems Biology, Harvard University, Cambridge, Massachusetts, USA^b; Département Micro-organismes, Génomes, Environnement, Equipe Adaptations et interactions microbiennes dans l'environnement, Université de Strasbourg, UMR 7156 UdS-CNRS, Strasbourg, France^c; Department of Chemical Engineering, University of Washington, Seattle, Washington, USA^d; IRD, Laboratoire des Symbioses Tropicales et Méditerranéennes, UMR 113, Montpellier, France^e; Department of Biochemistry and Molecular Biology, Wright State University, Dayton, Ohio, USA^f; Institute of Microbiology, ETH Zurich, Zurich, Switzerland^g; Bioscience Division, Los Alamos National Laboratory, Los Alamos, New Mexico, USA^h; DOE Joint Genome Institute, Walnut Creek, California, USAⁱ; Genomics Division, Lawrence Berkeley National Laboratory, Berkeley, California, USA^j; Laboratoire d'Analyses Bioinformatiques pour la Génomique et le Métabolisme (LABGeM), CEA-IG-Genoscope, Evry, France^k; Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA^l; and Pacific Northwest National Laboratory, Richland, Washington, USA^m

The complete and assembled genome sequences were determined for six strains of the alphaproteobacterial genus *Methylobacterium*, chosen for their key adaptations to different plant-associated niches and environmental constraints.

Genomic and metagenomic investigations have highlighted the prevalent role of methylotrophic microorganisms in a variety of marine, freshwater, and terrestrial environments (3–4, 6). These data have propelled new understanding of the molecular intricacies of microbial methylotrophic metabolism (1) and have sparked continued interest in their potential for biotechnological applications (15). In this work, the assembled complete genome sequences of six strains of the alphaproteobacterial genus *Methylobacterium* were determined. The selected strains were chosen for key characteristics, in terms of ecology, physiology, and metabolism (Table 1), in order to investigate how such adaptive features are reflected at the level of genome composition and architecture.

Genomes were sequenced at the Joint Genome Institute (JGI) using combinations of small to medium DNA libraries (3, 6, and 8 kb), as well as fosmid libraries (35 and 40 kb), with Sanger sequencing (7.3 to 9.6× coverage) completed with 454 pyrosequencing (20× coverage). All general aspects of library construction and sequencing can be found at http://www.jgi.doe.gov/sequencing/protocols/protos_production.html. Draft assemblies and quality assessment were obtained using the Phred/Phrap/Consed software package. Possible misassemblies were corrected with Dupfinisher (8), PCR amplification, and transposon bombing of bridging clones (Epicentre Biotechnologies, Madison, WI). Gaps between contigs were closed by editing in Consed, custom primer walking, and PCR amplification. A final assembly (7.5 to 10.5× coverage) was obtained for all 6 genomes (Table 1), and automatic annotation was performed using the JGI-Oak Ridge National Laboratory annotation pipeline (12). Additional automatic and manual sequence annotations, as well as comparative

genome analysis, were performed using the MicroScope platform at Genoscope (16).

The six *Methylobacterium* strains show significant variation in chromosome size and plasmid content (Table 1), and each possesses several conserved gene clusters known to be involved in methylotrophy in *Methylobacterium* (2, 18). Five of the strains possess conserved clusters of genes associated with photosynthesis, including genes encoding the light-harvesting complex and the reaction center, and genes involved in biosynthesis of bacteriochlorophyll and carotenoids. Further analyses of these six genomes will include comparisons to the two *Methylobacterium* genomes already reported (18), i.e., *M. extorquens* AM1, a major model strain in studies of methylotrophy (2) and genome evolution (5), and the dichloromethane-degrading strain *M. extorquens* DM4 (14). This will define both core- and strain-specific features of *Methylobacterium* strains and provide new insights into the metabolic flexibility of these facultative methylotrophs and into the modes of bacterial adaptation to specific ecological niches.

Nucleotide sequence accession numbers. GenBank accession numbers for all the chromosomes and plasmids sequenced in this study are shown in Table 1.

Received 6 June 2012 Accepted 13 June 2012

Address correspondence to Christopher J. Marx, cm Marx@oeb.harvard.edu, or Stéphane Vuilleumier, vuilleumier@unistra.fr.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.
doi:10.1128/JB.01009-12

TABLE 1 Characteristics of the six complete *Methylobacterium* genomes sequenced in this study

Organism	Key characteristic(s)	Genome analysis					GenBank accession no.	Reference		
		Size (Mb)	% GC	No. of <i>rrn</i> operons	No. of tRNAs	No. of CDS ^a				
<i>M. extorquens</i> strain PA1	<i>Arabidopsis thaliana</i> epiphyte	5.471	68.2	5	58	5,410	NC_010172	11		
<i>M. extorquens</i> strain CM4	Chloromethane degrader	5.778	68.2	5	61	3,112	NC_011757	13		
		0.380	66.3			388	NC_011758			
		0.023	63.9			44	NC_011760			
<i>M. extorquens</i> strain BJ001 ^b	<i>Populus deltoides</i> x <i>nigra</i> DN34 endophyte	5.800	69.4	5	58	6,017	NC_010725	17		
		0.025	64.9			30	NC_010727			
		0.023	66.8			31	NC_010721			
<i>M. radiotolerans</i> strain JCM 2831	Radioresistant strain	6.078	71.5	4	56	6,325	NC_010505	9		
		0.586	69.6			2	1		650	NC_010510
		0.047	62.5						66	NC_010509
		0.043	63.2				1		75	NC_010514
		0.038	63.7				1		60	NC_010517
		0.036	62.0						64	NC_010518
		0.028	61.0				1		45	NC_010502
		0.022	61.1						38	NC_010504
0.021	65.1			33	NC_010507					
<i>Methylobacterium</i> sp. strain 4-46	<i>Lotononis bainesi</i> nodulating, photosynthetic	7.659	71.6	6	63	8,337	NC_010511	7		
		0.058	65.1			108	NC_010373			
		0.020	59.2			34	NC_010374			
<i>M. nodulans</i> strain ORS 2060	Nonpigmented, nitrogen fixing, <i>Crotalaria</i> nodulating	7.772	68.9	7	71	8,879	NC_011894	10		
		0.488	65.9			2	630		NC_011892	
		0.458	65.7				609		NC_011887	
		0.040	64.2				84		NC_011893	
		0.038	61.6				66		NC_011895	
		0.020	61.4				30		NC_011888	
		0.013	60.5				16		NC_011889	
		0.010	67.2				14		NC_011890	

^a Number of annotated protein-coding sequences in MicroScope (16).

^b This strain, originally reported as *M. populi* strain BJ001 (17), was assigned to the species *M. extorquens* based on 16S rRNA gene identity (99.3%) and overall genome similarity with the four other sequenced *M. extorquens* strains (~80% identity over 75% of its genome sequence) (also see reference 18).

ACKNOWLEDGMENTS

This research resulted from an approved proposal (Marx_0165_051130) evaluated during the DOE-CSP-05 program supported by the Office of Biological and Environmental Research in the DOE Office of Science. The team of C.J.M. was also supported by an NSF grant (IOB-0612591), and the team of S.V. was also supported by a mobility grant from CNRS (USA mobility program). The work conducted by the U.S. Department of Energy Joint Genome Institute was supported by the Office of Science of the U.S. Department of Energy under contract DE-AC02-05CH11231.

We are grateful to the JGI personnel who participated in the sequencing, assembly, and automated annotation processes.

REFERENCES

- Anthony C. 2011. How half a century of research was required to understand bacterial growth on C1 and C2 compounds; the story of the serine cycle and the ethylmalonyl-CoA pathway. *Science Prog.* **94**:109–137.
- Chistoserdova L. 2011. Modularity of methylotrophy, revisited. *Environ. Microbiol.* **13**:2603–2622.
- Chistoserdova L. 2010. Recent progress and new challenges in metagenomics for biotechnology. *Biotechnol. Lett.* **32**:1351–1359.
- Chistoserdova L, Kaluzhnyaya MG, Lidstrom ME. 2009. The expanding world of methylotrophic metabolism. *Annu. Rev. Microbiol.* **63**:477–499.
- Chou H-H, Chiu H-C, Delaney NF, Segre D, Marx CJ. 2011. Diminishing returns epistasis among beneficial mutations decelerates adaptation. *Science* **332**:1190–1192.
- Delmotte N, et al. 2009. Community proteogenomics reveals insights into the physiology of phyllosphere bacteria. *Proc. Natl. Acad. Sci. U. S. A.* **106**:16428–16433.
- Fleischman D, Kramer DM. 1998. Photosynthetic rhizobia. *Biochim. Biophys. Acta* **1364**:17–36.
- Han C, Chain P. 2006. Finishing repetitive regions automatically with Dupfinisher, p 141–146. *In* Arabnia HR, Valafar H (ed), *Proceedings of the 2006 International Conference on Bioinformatics and Computational Biology*. CSREA Press, Las Vegas, NV.
- Ito H, Iizuka H. 1971. Part XIII: taxonomic studies on a radio-resistant *Pseudomonas*. *Agric. Biol. Chem.* **35**:1566–1571.
- Jourand P, et al. 2004. *Methylobacterium nodulans* sp. nov., for a group of aerobic, facultatively methylotrophic, legume root-nodule-forming and nitrogen-fixing bacteria. *Int. J. Syst. Evol. Microbiol.* **54**:2269–2273.
- Knief C, Frances L, Vorholt JA. 2010. Competitiveness of diverse *Methylobacterium* strains in the phyllosphere of *Arabidopsis thaliana* and identification of representative models, including *M. extorquens* PA1. *Microb. Ecol.* **60**:440–452.

12. Mavromatis K, et al. 2009. The DOE-JGI standard operating procedure for the annotations of microbial genomes. *Stand. Genomic Sci.* 1:63–67.
13. McDonald IR, Doronina NV, Trotsenko YA, McAnulla C, Murrell JC. 2001. *Hyphomicrobium chloromethanicum* sp. nov. and *Methylobacterium chloromethanicum* sp. nov., chloromethane-utilizing bacteria isolated from a polluted environment. *Int. J. Syst. Evol. Microbiol.* 51:119–122.
14. Muller EEL, et al. 2011. Functional genomics of dichloromethane utilization in *Methylobacterium extorquens* DM4. *Environ. Microbiol.* 13: 2518–2535.
15. Schrader J, et al. 2009. Methanol-based industrial biotechnology: current status and future perspectives of methylotrophic bacteria. *Trends Biotechnol.* 27:107–115.
16. Vallet D, et al. 2009. MicroScope: a platform for microbial genome annotation and comparative genomics. Database. doi:10.1093/database/bap021.
17. Van Aken B, Peres CM, Doty SL, Yoon JM, Schnoor JL. 2004. *Methylobacterium populi* sp. nov., a novel aerobic, pink-pigmented, facultatively methylotrophic, methane-utilizing bacterium isolated from poplar trees (*Populus deltoides* x *nigra* DN34). *Int. J. Syst. Evol. Microbiol.* 54:1191–1196.
18. Vuilleumier S, et al. 2009. *Methylobacterium* genome sequences: a reference blueprint to investigate microbial metabolism of C1 compounds from natural and industrial sources. *PLoS One* 4:e5584. doi:10.1371/journal.pone.0005584.